

NTUTH-97-04

July 1997

Perspectives on Quark Mass and Mixing Relations ^{*}

George Wei-Shu HOU [†]

Department of Physics, National Taiwan University,
Taipei, Taiwan 10764, R.O.C.

Abstract

Two down–up perspectives on quark mass–mixing relations are reviewed. The modified Fritzsch path relates V_{cb} and V_{ub} to M_u , but has trouble with the low $V_{cb} \simeq 0.04$ value. The modified Wolfenstein path focuses on the change in V_{ub} from λ^3 (ca. 1983) $\rightsquigarrow \lambda^4$ (ca. 1994). The relations $m_d/m_s \sim m_s/m_b \sim \delta \sim \lambda^2 \simeq |V_{cb}| \simeq |V_{us}|^2$ and $m_u/m_c \sim m_c/m_t \sim \delta^2 \sim \lambda^4 \sim |V_{ub}|$ suggest a clean separation of the origin of V_{KM} : $|V_{us}| \equiv \lambda$ and $|V_{cb}| \equiv \delta$ arise from M_d while $V_{ub} \equiv B\lambda^4 e^{-i\phi}$ comes from M_u . Five to six parameters might suffice for ten mass–mixing parameters, with δ seemingly the more sensible expansion parameter, while $\lambda \simeq \sqrt{m_d/m_s}$ is tied empirically to $(M_d)_{11} = 0$. The approximate relations suggest a near weak scale origin of flavor.

^{*} Talk presented at the Workshop on Masses and Mixings of Quarks and Leptons, University of Shizuoka, Japan, March 19 – 21, 1997. To appear in Proceedings.

[†] E-mail: wshou@phys.ntu.edu.tw.

Perspectives on Quark Mass and Mixing Relations

George Wei-Shu Hou

Department of Physics
National Taiwan University
Taipei, Taiwan 10764, R.O.C.

Abstract

Two down-up perspectives on quark mass-mixing relations are reviewed. The modified Fritzsch path relates V_{cb} and V_{ub} to M_u , but has trouble with the low $V_{cb} \simeq 0.04$ value. The modified Wolfenstein path focuses on the change in V_{ub} from λ^3 (ca. 1983) $\rightsquigarrow \lambda^4$ (ca. 1994). The relations $m_d/m_s \sim m_s/m_b \sim \delta \sim \lambda^2 \simeq |V_{cb}| \simeq |V_{us}|^2$ and $m_u/m_c \sim m_c/m_t \sim \delta^2 \sim \lambda^4 \sim |V_{ub}|$ suggest a clean separation of the origin of V_{KM} : $|V_{us}| \equiv \lambda$ and $|V_{cb}| \equiv \delta$ arise from M_d while $V_{ub} \equiv B\lambda^4 e^{-i\phi}$ comes from M_u . Five to six parameters might suffice for ten mass-mixing parameters, with δ seemingly the more sensible expansion parameter, while $\lambda \simeq \sqrt{m_d/m_s}$ is tied empirically to $(M_d)_{11} = 0$. The approximate relations suggest a near weak scale origin of flavor.

I. Introduction

Why am I slightly embarrassed in playing the mass-mixing game (MMG)? Though MMG as started by [1] Harald Fritzsch is fascinating, I always hesitated in writing something. I mean, *just how many papers can fit in a 3×3 matrix?*

This attitude is not unique to me, as the depressing *philosophical* debates with referees (when I fail the temptation...) often show:

- “Long on Words and Short on Substance. ... *ad hoc*.”
—— Indeed we are!
- “No Real Insight!”
“Papers should contain More than Assumptions!”
- “Void of Dynamics!”
↔ “Ansätze not based on Symmetry *do not go beyond Mere Numerology!*”
↔ “There is an ∞ of phenomenologically consistent and predictive Ansätze that Do Not Follow from Symmetry.”
↔ “**I do Not think they should be Published.**”

Well, does this ring a bell? Perhaps you even secretly agree when not sitting in the author’s pants? I would like to make the following,

STATEMENT

Flavor question very difficult!

— Available info limited. Just ask Mendeleev, Balmer, or Gell-Mann.

But fermion masses and mixing dominate # of free parameters in SM!

— Taking $m_\nu = 0$ and neglecting θ_{QCD} , one has $13/18 > 1/2$.

$\implies$ Reduction of Parameters Desirable!

Physics has always progressed on Idealization and establishing Empirical Rules or regularities, way before (and usually facilitating) Dynamical Explanations! So, I shall proceed with good conscience, since the *True Origins* of m_f , V_{KM} and CP violation are still *Unknown*. And, afterall, it’s the subject of this workshop.

There has been three paths in the mass-mixing game (as compared to much more serious “theory”, such as from topology, chaos, etc.):

1. Fritzsch [1] ($\sqrt{m_i/m_j}$): this will be our first theme [2].
2. Wolfenstein [3] (λ -expansion): this will be our second, main, theme [4].
3. Dimopoulos–Hall–Raby [5] (texture): will be commented on briefly.

Before we proceed, it is instructive to gain some perspective on the *scale* of m_f generation. We tend to think that “we” are normal, and “they” are abnormal, hence the top is *very heavy*, as we are made of u , d and e and need ν_e to get energy. Clearly $m_t \gg m_d$. Likewise, the W , Z , as well as H^0 bosons are also very heavy. But we should remember that all masses in SM are pinned to the v.e.v. scale, $v \sim 250$ GeV. We do not really know what happens at v.e.v. scale or beyond (call it “heaven”). From this perspective, as illustrated in Fig. 1, in fact the “heavy particles” appear to be *normal*, while we, you and I, seem to be made of *zero modes*. It is the latter that needs to be explained (*e.g.* pions as Goldstones), rather than the states with mass $\sim$ dynamical scale, in analogy with QED and QCD. This suggests that more states could appear at or above the v.e.v. scale. In this light, the mass-mixing hierarchy reflects a restoration of Decoupling (in spontaneously broken nonabelian gauge theories): Heavier Quarks do not play a major role in light quark physics (low energy phenomena). Fortified with this thought, let’s play game: Mass-Mixing relations without peeking above the v.e.v. scale.

One has the approximate relations $V_{ub}/V_{cb} \sim \sqrt{u/c}$, $V_{td}/V_{ts} \sim \sqrt{d/s}$, hence the smallness of V_{ub} compared to V_{cb} comes out quite naturally.

In 1978, bottom had just been discovered (in the form of Υ), but the top quark mass was not known at all. Fritzsche had hoped to predict m_t from V_{cb} , or *vice versa*, by tuning the cancellations for V_{cb} , which is highlighted in (3). The chain of colliders, PEP $\rightarrow$ PETRA $\rightarrow$ TRISTAN $\rightarrow$ LEP, were largely aimed at discovering the top quark, a hope vanquished by the ARGUS discovery of large B - $\bar{B}$ mixing. Today, the top quark mass has been measured at the Tevatron to be very, very heavy, completing the mass ratios

$$\frac{d}{s} \simeq \frac{1}{21} - \frac{1}{18} \gg \frac{u}{c} \sim \frac{1}{390} - \frac{1}{200}, \quad (4)$$

$$\frac{s}{b} \simeq \frac{1}{40} - \frac{1}{25} \gg \frac{c}{t} \sim \frac{1}{280}. \quad (5)$$

With such heavy top, does the Fritzsche Ansatz of (3) still work?

Critique: Main Problem from $V_{cb} \ll V_{us}$

With $V_{us} \simeq \sqrt{d/s}$ and $V_{cb} \simeq \sqrt{s/b} - \sqrt{c/t} e^{i\phi}$, since $\sqrt{s/b} \sim \sqrt{d/s}$, there are serious flaws even without m_t being extremely heavy. Below is a critique [2]:

- (a) Need $|\tau - \sigma| \lesssim 10^\circ - 15^\circ$ to allow for cancellation.
- (b) m_t *tuned* to cancel $\implies m_t \lesssim 60$ GeV. Is this arbitrary, or predictive?
- (c) $\sigma \approx \pm\pi/2 \longmapsto |V_{us}|^2 \sim d/s + u/c$
But combine with (a) $\implies$ CP violating phase *not* predicted.
- (d) And, because of (b): RULED OUT WHEN $m_t > M_W$.
Impossible to account for $V_{cb} \ll \sqrt{s/b} \sim V_{us} \nearrow$

Thus, the Fritzsche Ansatz was not very satisfactory in the first place.

He-Hou Variant: Relating $\tau_b \propto 1/m_t$

Dimopoulos once gave a talk at CERN and dubbed “He(e)-H(o)u did it.” Well, it really should be He[r]-Ho[u] phonetically. There is an interesting story about how it got started: He[r]-Ho[u] did it in Spring 1989 over some Löwenbraü (beer) sitting on Ludwigstrasse in München, not far from the Ludwig-Maximilians-Universität where Fritzsche is. Say we were inspired ...

We started by discussing the culprit: $\sqrt{s/b}$ was way too large for V_{cb} in (3). We said to ourselves: “*Just Dump it!*”, and went on with more beer ...

More sober the morning after, we made the

Postulate: No s - b or d - b mixing $\implies$ one rotates only in d - s plane.

With M_u unchanged from (1), our Ansatz amounted to

$$M_d = \begin{bmatrix} 0 & a & 0 \\ a & b & 0 \\ 0 & 0 & c \end{bmatrix}. \quad (6)$$

One immediately finds that (i) the phase τ could now be absorb into redefining b_L and b_R and (ii) one just sets s/b and $d/b \rightarrow 0$ in (3) and arrive at

$$V \simeq \begin{bmatrix} 1 & \sqrt{\frac{d}{s}} - \sqrt{\frac{u}{c}} e^{i\sigma} & -\sqrt{\frac{u}{t}} e^{i\sigma} \\ -\sqrt{\frac{d}{s}} + \sqrt{\frac{u}{c}} e^{-i\sigma} & 1 & \sqrt{\frac{c}{t}} \\ \sqrt{\frac{d}{s}} \sqrt{\frac{c}{t}} & -\sqrt{\frac{c}{t}} & 1 \end{bmatrix}, \quad (7)$$

which is considerably simpler. The salient features are:

- (a) $V_{cb} = \sqrt{c/t}$: We related Long τ_b (Smallness of V_{cb}) $\iff$ Heaviness of m_t without double fine-tuning as in the Fritzsch case.

Using QCD running only and ignoring the potentially more important weak corrections (running of λ_t), we found that

$$0.03 \lesssim |V_{cb}| \lesssim 0.06 \implies 750\text{GeV} \gtrsim m_t \gtrsim 200\text{GeV}. \quad (8)$$

This was a tantalizing result, but also indicative of the problem that we should not have ignored the running of λ_t .

- (b) $|V_{ub}/V_{cb}| = \sqrt{u/c}$ and $|V_{td}/V_{ts}| = \sqrt{d/s}$ become exact. Only V_{cb} is related to $1/m_t$, the rest of V_{KM} is expressed in terms of V_{cb} and ratios of masses of first two generations. The prediction of very small V_{ub} comes out naturally!

- (c) $\sigma \approx \pm\pi/2$ as in Fritzsch.

But, there is *only one single phase* $\longrightarrow$ CP violating phase determined.

- (d) Maximal CP: Strictly! That is, from

$$\begin{aligned} J_{\text{CP}} &\propto \text{Im}(V_{us}V_{ub}^*V_{cs}^*V_{cb}) \simeq \sqrt{\frac{m_d}{m_s}} \sqrt{\frac{m_u}{m_t}} \sqrt{\frac{m_c}{m_t}} \left(1 + \frac{m_u}{m_c}\right) \sin \sigma \\ &\simeq |V_{us}| |V_{ub}| |V_{cs}| |V_{cb}| \sin \sigma \cong (3-4) \times 10^{-5}, \end{aligned} \quad (9)$$

the CP phase is clearly maximal if $\sigma = \pm\pi/2$.

- (e) Wolfenstein expansion [3]: Take $\sin \theta \equiv \lambda$, $V_{cb} \sim \lambda^2$, and $V_{ub}/V_{cb} = \sqrt{u/c} \sim \lambda^2$, after some phase redefinitions, we find the Wolfenstein form

$$V = \begin{bmatrix} 1 & \lambda & \lambda^5 \mp i\lambda^4 \\ -\lambda & 1 & \lambda^2 \\ (1 \mp i\lambda)\lambda^3 & -\lambda^2 & 1 \end{bmatrix}. \quad (10)$$

The upshot is that we have a completely predictive Ansatz: *heavy top, maximal CP, and V_{KM} fixed in terms of $\sqrt{m_i/m_j}$ ratios!* The mass matrices of M_d in (6) and M_u in (1) were just (unkown to us!) the Georgi and Jarlskog (“texture”) form [9], hence the work anticipated the Top $\searrow$ Down approach of SO(10) SUSY-GUT models of Dimopoulos, Hall and Raby [5], which can be viewed as a high scale approach to remedy the over-heavy top.

As V_{cb} went down from ~ 0.06 in 1989 to ~ 0.04 by 1993, it became very difficult even for (7) to account for the smallness of V_{cb} . The problem remains severe even for the SO(10) based SUSY-GUT models.

III. “Wolfenstein” Path: from $|V_{cb}| \simeq |V_{us}|^2 \ll 1$

As we have mentioned the Wolfenstein parametrization in (10) already, it is useful to gain some historic perspective.

Then and Now: B Decay Data from 1983 $\rightsquigarrow$ 1994

It was discovered in 1983 that the B lifetime was much longer than expected. Together with the absence of $b \rightarrow u$ transitions, the new experimental values were

$$V_{cb} \approx 0.06, \quad |V_{ub}/V_{cb}| < 0.2, \quad (11)$$

indicating that $V_{ub}^2 \ll V_{cb}^2 \ll V_{us}^2$. Taking note of this, Wolfenstein suggested that [3] $\lambda \equiv V_{us} \simeq 0.22$ could be taken as an expansion parameter for V_{KM} , and proposed to parametrize the KM matrix as

$$V = \begin{bmatrix} 1 - \frac{1}{2}\lambda^2 & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \frac{1}{2}\lambda^2 & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{bmatrix}, \quad (12)$$

to order λ^3 , with $A \approx 5/4$ and $\rho^2 + \eta^2 < 1$ from (11). The parametrization has since become a reference standard [10] because of its value as a mnemonic device, especially in regards the popular unitarity triangle representation for CP violation,

Fourteen years has elapsed since Wolfenstein's proposal of (12). With the advent of CLEO II data and the development of HQET, the values for V_{cb} and $|V_{ub}/V_{cb}|$ have been consistently dropping in the past few years [10], stabilizing more or less by 1994. The current values are

$$|V_{cb}| = 0.040 \pm 0.005, \quad |V_{ub}/V_{cb}| = 0.08 \pm 0.02. \quad (13)$$

Thus, we now have

$$A = 0.8 \pm 0.1, \quad (14)$$

which is *down by 1/3* compared to ten years ago. But the drop in $|V_{ub}|$ is more dramatic: **a factor of 4 down** from that of (11), giving

$$|V_{ub}| = 0.0032 \pm 0.0009. \quad (15)$$

A factor of 1/4 corresponds to $\sim \lambda$ suppression. Noting that $\lambda \cong 0.2205$ hence $\lambda^2 \cong 0.0486$, $\lambda^3 \cong 0.0107$ and $\lambda^4 \cong 0.0024$, we can now define [4]

$$V_{ub} \equiv A\lambda^4(\rho' - i\eta') \equiv B\lambda^4 e^{-i\phi}, \quad (16)$$

where $\rho \equiv \rho'\lambda$, $\eta \equiv \eta'\lambda$ (i.e. $d\rho/d\lambda = \rho'$ as a memory device) and $\phi \equiv \tan^{-1} \eta'/\rho'$. Numerically,

$$B \equiv A\sqrt{\rho'^2 + \eta'^2} = 1.3 \pm 0.5, \quad (17)$$

and interestingly, $AB \sim 1$. We note that the original Wolfenstein parametrization is still better suited as a mnemonic device (see Fig. 2), but the change in powers of λ for V_{ub} might have some significance.




Figure 2: Unitarity triangle illustrating the change $V_{ub} : \lambda^3 \rightsquigarrow \lambda^4$.

Whether this is mere numerology or not depends on how you take the original Wolfenstein paper. The original proposal emphasized a possible λ -expansion [3], a point which was largely lost because of its great success as a mnemonic device. Afterall, Wolfenstein freely admitted that $\lambda \simeq 0.22$ is not so small as an expansion parameter, making the order of the series, if any, ambiguous.

If one takes the series expansion idea seriously [4], however, the change in $V_{ub} : \lambda^3 \rightsquigarrow \lambda^4$ amounts to a change in order of λ from *odd* $\rightsquigarrow$ *even*. This may have profound implications for possible underlying dynamics that relates masses and mixing angles. With the recent measurement of $m_t(m_t) \cong 175$ GeV, it is intriguing to note that, as shown in Eqs. (4) and (5) we now have

$$\frac{m_d}{m_s} \sim \frac{m_s}{m_b} \sim \lambda^2 \gg \frac{m_u}{m_c} \sim \frac{m_c}{m_t} \sim \lambda^4. \quad (18)$$

The relations suggest that V_{KM} comes mostly from down quark sector, while the up-type quark sector contributes only at higher order in λ . Let us therefore explore along these lines in more detail.

“Gestalt” Switch: Hint from $\boxed{V_{ub} : \lambda^3 \longmapsto \lambda^4}$

Combining Fritzsch with Wolfenstein, we have the old empirical relation

$$V_{us} = \lambda \simeq \sqrt{\frac{m_d}{m_s}}. \quad (19)$$

With the falling V_{cb} , several authors have noted [11, 12] that the relation

$$V_{cb} \equiv A\lambda^2 \simeq \frac{m_s}{m_b}, \quad (20)$$

now holds rather well. It appears then that *the KM matrix V is mostly due to the down type sector, while eqs. (16) and (18) suggest that perhaps V_{ub} — hence CP violation — originates from the up type sector*. We thus reach the Proposition:

$$D \sim V_{\text{KM}}. \quad (21)$$

Let us see how far we can go with this thought.

For simplicity, let us assume hermitian mass matrices (this limits the form of the possible underlying dynamics). Ignoring the u -type quark mass matrix to first approximation since its mass ratios are subdominant, we can reconstruct the down type quark mass matrix by rotating back with the KM matrix, that is

$$M_d \cong V \begin{bmatrix} -m_d & 0 & 0 \\ 0 & m_s & 0 \\ 0 & 0 & m_b \end{bmatrix} V^\dagger. \quad (22)$$

- Approximate “Texture Zero”: $\boxed{(M_d)_{11} = 0}$

The empirical relation of (19) can be maintained IF [13] $(M_d)_{11} = 0$ in (22), which is nothing but the old texture zero of Weinberg. It holds approximately true numerically, namely

$$(M_d)_{11} = -m_d|V_{ud}|^2 + m_s|V_{us}|^2 + m_b|V_{ub}|^2 \cong 0, \quad (23)$$

as can be most easily checked by making a series expansion in λ .

[In 1991, one still had $V_{ub} \sim \lambda^3$. Ma attempted, therefore, at a second zero, $(M_d M_d^\dagger)_{12} \simeq 0$, suggesting a second relation [13] $m_s^2/m_b^2 = -V_{ub}V_{cb}/V_{us}$. Both were of order λ^4 in 1991. With (13) hence (16), however, the righthand side is now of order λ^5 , and the relation no longer holds.]

- Idealize: *Least Number of Parameters*

To maintain the relation of (20), without loss of generality, let us redefine quark fields to make M_d real symmetric. This relegates all CP violating effects to the up-type sector, *i.e.* V_{ub} is from M_u . We therefore idealize our proposition of (21) and

$$\textbf{Postulate: } \boxed{D_L \equiv V_{\text{KM}}(V_{ub} = 0)}, \quad (24)$$

which corresponds to the *least number of parameters needed to account for both d-type quark masses and V , with $V_{ub} = 0$* . There are just two parameters, λ and A , or, we define $\lambda \equiv V_{us}$, $\delta \equiv V_{cb} \cong A\lambda^2$,

$$D_L = \begin{bmatrix} \sqrt{1-\lambda^2} & \lambda & \boxed{0} \\ -\lambda\sqrt{1-\delta^2} & \sqrt{1-\lambda^2}\sqrt{1-\delta^2} & \delta \\ \lambda\delta & -\sqrt{1-\lambda^2}\delta & \sqrt{1-\delta^2} \end{bmatrix}. \quad (25)$$

With $(D_L)_{13} = 0$ by fiat, Ma’s observation is now reformulated as

$$\begin{aligned} (M_d)_{11} &= -m_d(D_L)_{11}^2 + m_s(D_L)_{12}^2 \\ &= -m_d(1-\lambda^2) + m_s\lambda^2 = 0. \end{aligned} \quad (26)$$

Enforcing this “texture” zero strictly gives

$$\frac{m_d}{m_s} = \frac{\lambda^2}{1-\lambda^2}, \quad (27)$$

to all orders in λ , leading to the relation

$$V_{us} \equiv \lambda = \sqrt{\frac{m_d}{m_s + m_d}} \cong s_{12}, \quad (28)$$

or $\tan \theta_{12} = \sqrt{m_d/m_s}$. We see that the idealization (25) of our proposition (21) leads to a refinement of Eq. (19).

- Second “Texture Zero”: $\boxed{(M_d)_{13} = 0}$

Coming back to relation (20), we note that it implies $m_s : m_b = A\lambda^2 : 1$, while the all orders relation (27) gives $m_d : m_s = \lambda^2 : 1 - \lambda^2$. Taken together they suggest that

$$-m_d : m_s : m_b = -A\lambda^4 : A\lambda^2 - A\lambda^4 : 1, \quad (29)$$

Of course, we are not sure of corrections to the relation (20) since m_s , as well as m_b , are less precisely known as V_{cb} . But if we take the ratio (29) as is and idealize, we get an additional texture zero (M_d is real symmetric). Define $\hat{M}_d = M_d/m_b$, multiplying out (22), we find

$$\hat{M}_d = \begin{bmatrix} 0 & A\lambda^3 & 0 \\ A\lambda^3 & A\lambda^2 + (A-2)A\lambda^4 & A\lambda^2 - A^2\lambda^4 \\ 0 & A\lambda^2 - A^2\lambda^4 & 1 - A^2\lambda^4 \end{bmatrix}. \quad (30)$$

The texture zero of $(M_d)_{13} = 0$ implies the relation

$$V_{cb} \equiv A\lambda^2 = \frac{m_s + m_d}{m_b} \cong s_{23}, \quad (31)$$

which modifies Eq. (20) slightly, but is less convincing than (28).

Thus, with the three parameters m_b (scale parameter), A (~ 1) and λ , one can account for the 5 parameters m_d , m_s , m_b and $|V_{us}|$, $|V_{cb}|$, resulting in two relations, Eqs. (28) and (31).

Up-type Sector: CP Violation, m_u , and V_{ub}

Let us extend the previous arguments to the up-type sector. With the postulate of (24) that D_L accounts for V_{KM} up to $V_{ub} = 0$, we simply have

$$U_L \equiv D_L V^\dagger \simeq \begin{bmatrix} 1 & 0 & -B\lambda^4 e^{-i\phi} \\ 0 & 1 & 0 \\ B\lambda^4 e^{i\phi} & 0 & 1 \end{bmatrix} + \mathcal{O}(\lambda^6). \quad (32)$$

Taking M_u to be hermitian also, we have, analogous to Eq. (22),

$$\hat{M}_u = U_L \begin{bmatrix} -\hat{m}_u & 0 & 0 \\ 0 & \hat{m}_c & 0 \\ 0 & 0 & \hat{m}_t \end{bmatrix} U_L^\dagger, \quad (33)$$

where we normalize with respect to m_t , *i.e.* $\hat{m}_t = 1$. Noting the approximate orders $m_t : m_c : m_u \approx 1 : \lambda^4 : \lambda^8$ from (18), we multiply out (33) and get

$$\hat{M}_u = \begin{bmatrix} -\hat{m}_u + B^2\lambda^8 & 0 & -B\lambda^4 e^{-i\phi} \\ 0 & \hat{m}_c & 0 \\ -B\lambda^4 e^{i\phi} & 0 & 1 \end{bmatrix} + \mathcal{O}(\lambda^{10}). \quad (34)$$

We can now proceed to idealize.

- “Texture Zero”: $(M_u)_{11} = 0$ analogy

The $(\hat{M}_u)_{11}$ term is at order λ^8 and can be removed by invoking a texture zero condition analogous to but weaker than (26),

$$(M_u)_{11} = -m_u |U_L|_{11}^2 + m_t |U_L|_{13}^2 = 0, \quad (35)$$

which holds at least up to order λ^{12} . This leads to the relation

$$|V_{ub}| \equiv B\lambda^4 \cong \sqrt{\frac{m_u}{m_t}}, \quad (36)$$

hence *both* V_{ub} and m_u are generated via diagonalizing the u - t mixing element, analogous to the original $\sin \theta_C \simeq \sqrt{d/s}$ relation.

- Reduction of Parameters: *Least Number*

Eq. (18) suggest that we could write $\hat{m}_c = C\lambda^4$ with $C \simeq 1$, hence

$$\hat{M}_u \cong \begin{bmatrix} 0 & 0 & -B\lambda^4 e^{-i\phi} \\ 0 & C\lambda^4 & 0 \\ -B\lambda^4 e^{i\phi} & 0 & 1 \end{bmatrix}, \quad (37)$$

where the 11 element is at $\mathcal{O}(\lambda^{12})$, while the other 0's are at $\mathcal{O}(\lambda^{10})$. Since both C and B are ~ 1 , setting $C = B$ reduces the number of parameters, leading to a second, “geometric” relation

$$\frac{m_u}{m_c} = \frac{m_c}{m_t} = B\lambda^4. \quad (38)$$

We are now really deep into numerology. Since from (14) and (17) we have $A \simeq 1 - \lambda$ and $B \simeq 1 + \lambda$, it may well be that $B = 1/A$ or 1 in Nature, such that $m_u/m_c = m_c/m_t = |V_{ub}| = \lambda^4$ or $A^{-1}\lambda^4$, and just *two* additional parameters, m_t and ϕ , *might* account for the remaining *five* parameters m_u , m_c , m_t and $V_{ub} = |V_{ub}|e^{-i\phi}$. Given the uncertainties in mixing, and especially in the *lighter* quark masses [10], these results are not inconsistent with data! Note that the CP violating phase $\phi = \tan^{-1}(\eta'/\rho')$ is a free parameter, and does not arise from mass ratios.

- Naturalness of V_{ub} (and CP violation) from M_u

Eqs. (16–20) imply that $m_d/m_b = A\lambda^4 \lesssim m_u/m_c \sim m_c/m_t \sim |V_{ub}| \sim B\lambda^4$, so we could arrange to have V_{ub} to arise from $\hat{M}_d$ while $\hat{M}_u \simeq \text{diag}(-\lambda^8, \lambda^4, 1)$ to be already diagonal. This corresponds to taking the off-diagonal piece from U_L in (32) and placing it in D_L . However, this is *unnatural* since if V_{ub} comes from M_d and starts at $\mathcal{O}(m_d/m_b)$, then

$$\begin{cases} V_{ub} \propto m_d/m_b \\ m_t \text{ arbitray and large} \end{cases} \implies \varepsilon_K \text{ easily fails!}$$

It is therefore important, as in (32) or (37), to couple

Smallness of CP Violation Effects $\iff$ *Heaviness of the Top Quark*,

like in (38), where $|V_{ub}| \sim \sqrt{u/t} \sim c/t$, leading to

$$\varepsilon_K \propto \text{Im}(V_{ub})^2 m_t^2, \quad (39)$$

which would be stable against the rise of m_t . The placement of CP phase in $(M_u)_{13}$ may therefore be rooted in dynamics.

Discussion

- Since we are just applying the λ -expansion, the phenomenology is the same as in SM. In particular, the invariant area of the unitarity triangle

$$J_{CP} = \text{Im}(V_{us}V_{ub}^*V_{cs}^*V_{cb}) \simeq A^2\eta'\lambda^7 \lesssim 3 \times 10^{-5}, \quad (40)$$

is of order λ^7 . Though it is still convenient to use ρ and η of Wolfenstein [10] for the unitarity triangle, but with ρ , $\eta = \rho'\lambda$, $\eta'\lambda$, and ρ' , $\eta' \sim 1$ the unitarity triangle appears a bit squashed (see Fig. 2). The CP phase angle $\phi \equiv \tan^{-1} \eta'/\rho'$ remains a free parameter.

- δ -expansion *vs.* λ -expansion

It would be appealing if some symmetry or dynamical mechanism underlies the possible reduction of 2 to 5 parameters from the 10 quark masses and mixing angles. Discrete symmetries à la Ma [13] can be constructed by adding extra Higgs doublets. However, these usually do not add insight to the λ^n power behavior for mixing angles and mass ratios. Note that eqs. (30) and (37) suggest an expansion in even powers of λ , *except* for $(\hat{M}_d)_{12} = (\hat{M}_d)_{21} \cong \lambda^3$. This is because, with V_{ub} changed from order λ^3 to order λ^4 , the only term odd in λ is just $|V_{us}| = |V_{cd}|$ itself. Defining as before $\delta \equiv V_{cb} \equiv A\lambda^2$, we find, to leading order in δ ,

$$\hat{M}_d \cong \begin{bmatrix} 0 & \delta^{3/2} & 0 \\ \delta^{3/2} & \delta & \delta \\ 0 & \delta & 1 \end{bmatrix}, \quad \hat{M}_u \simeq \begin{bmatrix} 0 & 0 & -\delta^2 e^{-i\phi} \\ 0 & \delta^2 & 0 \\ -\delta^2 e^{i\phi} & 0 & 1 \end{bmatrix}. \quad (41)$$

So perhaps $\delta \simeq 1/20 - 1/30$ is the actual expansion parameter. The Wolfenstein's $\lambda = \sqrt{\delta/A}$ is empirically tied to $(\hat{M}_d)_{11} = 0$ [13], but one needs to understand the relatively large ratio m_d/m_s .

- The 1991 ansatz of Ng and Ng [12] also starts with the approximate relations of (18), but they impose $(M_d)_{32} = 0$, which we find unnecessary. The 1992 Ansatz of Giudice [14] starts from SUSY GUTS (Georgi–Jarlskog texture), and relates M_d and M_e with *ad hoc* assumptions, and arbitrarily sets $(M_d)_{23} = 2(M_d)_{22}$. Both of these works do not utilize the λ -expansion.

- General Texture Analysis

In 1993 Ramond, Roberts and Ross [15] made a systematic extension of the Dimopoulos–Hall–Raby approach. They studied all the possible *texture zeros* (which implies mass-mixing relations) of symmetric or hermitian M_u and M_d matrices *at SUSY GUT scale*, then evolved them down from Λ_{GUT} to Λ_{Weak} . They find 5 textures, the third of which is close to ours (see their Tables 1 and 2). They do make a λ -expansion in a special way, with $\sqrt{\rho^2 + \eta^2} \sim \lambda$. They view λ as possibly emerging from ratios of v.e.v.'s. But because the textures exist at GUT scale, they have the same old generic problem of DHR, namely it is difficult to bring V_{cb} below 0.045 (see their Table 3).

- Weak Scale Mass Generation?

In comparison, our Ansatz works BELOW WEAK SCALE, without any attempt to evolve upwards. We find (41) to be suggestive of an underlying

radiative mechanism, perhaps not far above the electroweak scale [16]. For example, the parameter $\delta \equiv A\lambda^2$ defined in eq. (24) could arise from loop processes at order $f^2/16\pi^2 = \alpha_f/4\pi$, where α_f comes from some underlying flavor dynamics. It need not have a high scale origin such as SUSY and/or GUTS [15]. We believe that the mass and mixing hierarchies, with correlations as exemplified in (41), cannot be just an accident.

IV. Conclusion

We have covered two generic approaches to the quark mass-mixing problem. Because of the scarcity of data, this is by far not the final word on this difficult but fascinating subject. However, we also do not pretend that we know what happens at way beyond the weak scale, nor demand that physics far beyond the weak scale is necessary to explain the flavor problem.

We think that the traditional Fritzsch path eventually runs into the difficulty of smallness of V_{cb} , and remedies such as (20) deviate drastically from the old Weinberg Ansatz for the Cabibbo angle. So, instead, we have emphasized and taken up the series expansion approach of Wolfenstein. Let us summarize the salient features:

- In 1983, Wolfenstein parametrized $V_{ub} \sim \lambda^3$. However, recent data give $V_{ub} \sim 0.003$, $V_{cb} \sim 0.04$ and $m_t \sim 175$ GeV. This leads to Eq. (16), that is

$$V_{ub} = A\lambda^4(\rho' - i\eta') \equiv B\lambda^4 e^{-i\phi}$$

where $\rho' \equiv d\rho/d\lambda$ and $B \simeq 1.3$ vs. $A \simeq 0.8$. So, from 1983 to 1993-1994, we have the change

$$V_{ub} : \lambda^3 \longmapsto \lambda^4.$$

This is the key starting point to revisit Wolfenstein's approach.

- Turning to mass ratios, from Eq. (18), that is

$$\frac{m_d}{m_s} \sim \frac{m_s}{m_b} \sim \lambda^2 \gg \frac{m_u}{m_c} \sim \frac{m_c}{m_t} \sim \lambda^4.$$

the mass ratios in the up-type sector are of higher order nature. This in turn suggests that V_{KM} originates mostly from down-type sector, Eq. (24), while V_{ub} arises from up-type quarks, Eq. (32). Identifying $\delta \equiv A\lambda^2 \sim 0.04$ as the more appealing expansion parameter, we have

$$\left\{ \begin{array}{l} D_L \equiv V_{\text{KM}}(V_{ub} = 0) \quad \Longleftarrow \quad \hat{M}_d \cong \begin{bmatrix} 0 & \delta^{3/2} & 0 \\ \delta^{3/2} & \delta & \delta \\ 0 & \delta & 1 \end{bmatrix}, \\ U_L \equiv D_L V_{\text{KM}}^\dagger \quad \Longleftarrow \quad \hat{M}_u \simeq \begin{bmatrix} 0 & 0 & -\delta^2 e^{-i\phi} \\ 0 & \delta^2 & 0 \\ -\delta^2 e^{i\phi} & 0 & 1 \end{bmatrix}. \end{array} \right.$$

Wolfenstein's parameter $\lambda \sim \sqrt{\delta}$ is the more puzzling, arising empirically from the famous Weinberg texture zero of $(M_d)_{11} = 0$, Eq. (26), which holds very well with 3 generation masses and mixings.

- Demanding the *least number of parameters*, which is related in spirit but not equivalent to finding texture zeros, we suggest the following approximate relations (Eqs. (28), (31) and (36)):

$$\begin{aligned} \lambda &= \sqrt{\frac{m_d}{m_s + m_d}} = |V_{us}|, \\ A\lambda^2 &= \frac{m_s + m_d}{m_b} = |V_{cb}|, \\ B\lambda^4 &= \sqrt{\frac{m_u}{m_t}} = |V_{ub}|, \quad \left(\frac{u}{c} = \frac{c}{t} = B\lambda^4\right), \end{aligned}$$

by idealizing the input equations (16)–(20).

- Note that $A \sim 1 \sim B$, but the smallness of m_b/m_t is not explained. Since $m_b/m_t = (m_b/m_c)(m_c/m_t) \sim (m_b/m_c)\lambda^4 \sim A\lambda^3$, perhaps both A and the odd power λ expansion are related to m_b generation from a heavy top.
- The placement of CP phase in $(M_u)_{13}$ may be rooted in dynamics, since it naturally *couples the smallness of CP violation effects to the existence and heaviness of the top quark*.
- Since this “numerology” seems to work well with present, low energy data, it suggests *WEAK SCALE ORIGIN!*

Acknowledgments. The works reported here were done in collaboration with Xiao-Gang He and Gwo-Guang Wong, respectively.

References

- [1] H. Fritzsch, Phys. Lett. **70B**, 436 (1977).
- [2] X. G. He and W. S. Hou, Phys. Rev. **D 41**, 1517 (1990).
- [3] L. Wolfenstein, Phys. Rev. Lett. **51**, 1945 (1983).
- [4] W. S. Hou and G.G. Wong, Phys. Rev. **D 52**, 5269 (1995).
- [5] S. Dimopoulos, L. J. Hall, and S. Raby, Phys. Rev. Lett. **68**, 1984 (1991).
- [6] R. Gatto, G. Sartori, and M. Tonin, Phys. Lett. **28B**, 128 (1968); N. Cabibbo and L. Maiani, *ibid.* **28B**, 131 (1968).
- [7] S. Weinberg, Transactions of the New York Academy of Sciences, Series II, Vol. 38, 185 (1977).
- [8] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [9] H. Georgi and C. Jarlskog, Phys. Lett. **86B**, 297 (1979).
- [10] Review of Particle Properties, Phys. Rev. **D 50**, 1173 (1994).
- [11] K. S. Babu and R. N. Mohapatra, Phys. Rev. Lett. **64**, 2747 (1990).
- [12] D. Ng and Y. J. Ng, Mod. Phys. Lett. **A6**, 2243 (1991).
- [13] E. Ma, Phys. Rev. **D 43**, R2761 (1991).
- [14] G. F. Giudice, Mod. Phys. Lett. **A7**, 2429 (1992).
- [15] P. Ramond, R. G. Roberts, G. G. Ross, Nucl. Phys. **B 406**, 19 (1993).
- [16] For discussion of the lepton sector, see G. G. Wong and W. S. Hou, Phys. Rev. **D 50**, R2962 (1994). We ventured into quark mass-mixing numerology as an effort to extend from leptons to quarks.